

Bharati Vidyapeeth's College of Pharmacy

Sector-8, CBD Belapur, Navi Mumbai. 400614

SAMPLE QUESTIONS (MCQs)


Final Year B. Pharm. Sem-VII

PHARMACEUTICAL CHEMISTRY II - CBCS SYLLABUS

Q.1 All drugs are alkylating agents except..

- a) Busulfan
- b) Cyclophosphamide
- c) Chloramducil
- d) Methotrexate

Q.2 IUPAC name of this drug is..



- a) 2-amino-3-{4-[bis(2-chloroethyl)amino]phenyl} propanoic acid
- b) 1-amino-3-{4-[bis(2-chloroethyl)amino]phenyl} propanoic acid
- c) 2-amino-3-{4-[bis(2-chloroethyl)amino]phenyl} ethanoic acid
- d) 2-amino-3-{4-[(2-chloroethyl)amino]phenyl} propanoic acid

Q.3 Active metabolite of 6-Thioguanine is ..

- a) 6-Thioinosinic acid
- b) 6-Thioguanilyc acid
- c) 4-Thioguanilyc acid
- d) 4-Thioinosinic acid

Q.4 Which starting material is used for the synthesis of Chlorambucil?

- a) Benzanilide and phthalic anhydride
- b) Benzanilide and succinic anhydride
- c) Acetanilide and succinic anhydride
- d) Acetanilide and phthalic anhydride


Q.5 Select 1st generation antihistaminic agent.

- a) Cetirizine
- b) Levocetirizine
- c) Promethazine
- d) Fexofenadine

Q.6 Which sentence is wrong with reference to SAR of 1st generation antihistaminics?

- a) Two aromatic rings are essential connected to central carbon, nitrogen or carbon-oxygen
- b) Spacer between amino terminal amino nitrogen and central carbon atom should be 2-3 carbon atoms
- c) Amine is substituted with small alkyl groups.
- d) Potency of terminal amino nitrogen is –primary > secondary > tertiary

Q.7 What is the name of this drug?



- a) Omeprazole
- b) Rabeprazole
- c) Pentoprazole
- d) Ranitidine

Q.8is an oxidative metabolite of Terfenadine

- a) Fexofenadine
- b) Loratidine
- c) Cetirizine
- d) Ebastine

Q.9 Which is a second generation sulfonylurea?

- a) Tolbutamide
- b) Glipizide
- c) Chlorpropamide
- d) Acetohexamide

Q.10 Thiazolidinediones are...

- a) PPAR-gamma antagonist
- b) GLP-1 agonist
- c) PPAR-gamma agonist
- d) -Glucosidase inhibitors

Q.11 Select Nucleoside reverse transcriptase inhibitor.


- a) Delaviridine
- b) Nevirapine
- c) Efavirenz

d) Lamivudine

Q.12 What is the structural difference between Acyclovir and Ganciclovir?

- a) Ganciclovir contains an extra hydroxymethylene group
- b) Acyclovir contains an extra hydroxymethylene group
- c) Ganciclovir contains an extra methyl group
- d) Ganciclovir contains an extra methoxy group

Q.13 What is the IUPAC name of the following drug?




- a) 5-(acetoxymethyl)-3-[2-(dimethylamino) ethyl]-2-(4-methoxyphenyl) -1,5-benzothiazepin-4(5H)-one
- b) 3-(acetoxymethyl)-5-[2-(dimethylamino) methyl]-2-(4-methoxyphenyl) -1,4-benzothiazepin-4(5H)-one
- c) 3-(acetoxymethyl)-5-[2-(dimethylamino) ethyl]-2-(4-methoxyphenyl) -1,5-benzothiazepin-4(5H)-one
- d) 3-(acetoxymethyl)-5-[2-(dimethylamino) ethyl]-2-(4-methoxyphenyl) -1,5-benzothiazepin-4(5H)-one

Q. 14 Losartan can be classified as

- a) Angiotensin Converting enzyme inhibitor
- b) Angiotensin II receptor blocker
- c) Renin inhibitor
- d) K channel opener

Q. 15 Following starting materials will be used for synthesis of which drug?




- a) Prazosin
- b) Spironolactone
- c) Sotalol
- d) Captopril

Q. 16 Which of the following drug is nonselective β blocker

- a) Propranolol
- b) Acebutolol
- c) Esmolol
- d) Terazosin

Q. 17 with respect to the following general structure which is the appropriate statement?



- a) R' and R'' must be identical for optimum activity
- b) R must be at ortho or meta position for optimum activity
- c) R must be at para position for optimum activity
- d) conversion of 1,4-dihydro pyridine ring to pyridine ring increases activity


Q. 18 Diuretic containing phenoxy acetic acid scaffold is

- a) Acetazolamide
- b) Amiloride
- c) Mannitol
- d) ethacrynic acid

Q. 19 Which of the following is a fibrate prodrug?

- a) Gemfibrozil
- b) Ciprofibrate
- c) Clofibrate
- d) Carvedilol

Q. 20 Following antiarrhythmic drug will be classified under which class?



- a) Class I A
- b) Class I B
- c) Class I C
- d) Class II

Q. 21 Which of the following two drugs require hydrolysis for metabolic activation?

- a) Lovastatin and Simvastatin
- b) Lovastatin and Atorvastatin
- c) Simvastatin and Rosuvastatin
- d) Atorvastatin and Rosuvastatin

Q. 22 Which of the following Anticoagulant is Indanedione derivative?

- a) Aspirin
- b) Cilostazol
- c) Abciximab
- d) Phenindione

Q. 23 Which of the following anti-arrhythmic drug prevents peripheral conversion of T4 to T3?

- a) Propafenone
- b) Amiodarone
- c) Mexiletine
- d) Lidocaine

Q. 24 Which of the following drugs have been developed from benzyl succinic acid?

- a) Captopril
- b) Candesartan
- c) Aliskiren
- d) Eplerenone

Q. 25 Which of the following is irreversible COX -1 inhibitor

- a) Indobufen
- b) Aspirin
- c) Clopidogrel
- d) Fibrinogen